



Philip, N., Bandara, H. M. H. N., Leishman, S. J., & Walsh, L. J. (2019). Effect of polyphenol-rich cranberry extracts on cariogenic biofilm properties and microbial composition of polymicrobial biofilms. *Archives of Oral Biology*, 102, 1-6.
<https://doi.org/10.1016/j.archoralbio.2019.03.026>

Peer reviewed version

License (if available):
CC BY-NC-ND

Link to published version (if available):
[10.1016/j.archoralbio.2019.03.026](https://doi.org/10.1016/j.archoralbio.2019.03.026)

[Link to publication record in Explore Bristol Research](#)
PDF-document

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Table 1. Inhibitory effects of cranberry extract on biomass and lactic acid production of polymicrobial biofilms

Treatment Group	Biofilm biomass (% change) (OD ₅₇₀)	Lactic acid production (% change) (mM/l)
Cranberry	0.089 ± 0.012 (-38)*	6.2 ± 1.9 (-44)*
Control	0.144 ± 0.022	11.2 ± 3.8

Biomass represents mean crystal violet absorbance values ± standard deviation (SD) at optical density (OD) of 570 nm. Lactic acid production of biofilms is in mM/l (mean ± SD). Data for both assays is obtained from three independent triplicate experiments (n = 9). Values in brackets represent percentage reduction of biofilm biomass/acid production compared to control. * indicates significant difference compared to vehicle control using independent sample-t-test/Mann-Whitney *U*-test at confidence interval of 95%

Table 2. Quantitative analysis of biovolumes of polymicrobial biofilms from confocal images

Treatment Group	EPS biovolume ($\mu\text{m}^3/\mu\text{m}^2$)	Microbial biovolume ($\mu\text{m}^3/\mu\text{m}^2$)	Total biofilm biovolume ($\mu\text{m}^3/\mu\text{m}^2$)	EPS/Microbial ratio
Cranberry	1.14 \pm 0.15*	2.05 \pm 0.55*	3.19 \pm 0.63*	0.59 \pm 0.14*
Control	2.64 \pm 0.56	3.82 \pm 0.73	6.46 \pm 1.19	0.70 \pm 0.12

Biovolume data was quantified using IMARIS and represents mean values \pm SD (n=15) from 3 independent experiments.

* indicates significant difference ($p < 0.05$) in cranberry-treated biofilms compared to control using independent sample t-test.